

PRE-SHOW

- Practice good biosecurity on your farm
 - Do not share equipment or other supplies with other flocks
 - Limit visitors to your flock and avoid visits to poultry facilities or other backyard flocks
 - Wear designated, clean, disinfected footwear when caring for your flock to reduce the risk of contamination with wild bird feces and other contaminants
 - Do not allow your birds to mingle with wild birds
 - House indoors or in a screened in area
 - Practice pest and rodent control
 - Keep area clean
 - Keep feed covered and clean up spills
- Monitor your birds before the show, especially those you plan to take
 - Monitor for at least 14 days prior to show
 - Watch for changes in food and water consumption which can be an initial indicator of illness
 - Monitor for any potential clinical signs including respiratory signs and lethargy
 - Isolate sick birds from the rest of your flock & tend to them last
 - Report any clinic signs noted to your veterinarian
 - REMEMBER ONLY HEALHTY BIRDS CAN BE BROUGHT TO A SHOW, IF ANY BIRDS IN THE FLOCK ARE SICK, DO NOT ATTEND THE SHOW





- Keep your cages and feed/water supplies clean. Clean and disinfect your cages after the show.
- Limit contact with any other flocks or wild birds at the show
 - Keep species separate; do not mingle waterfowl and other poultry species
- Wash hands often and especially after handling birds
- Monitor all birds on site for clinical signs
 - If you see something in your flock or any other, notify the person in charge so that birds can be removed and returned home or isolated ASAP



POST-SHOW

- Isolate the birds that were at the show for 30 days before returning to flock
 - Monitor for any signs of illness
 - Tend to these birds after caring for the rest of your flock
- Clean and disinfect all cages and equipment that were at the show before using at home
- Clean and disinfect footwear worn at show
- It is recommended that exhibitors wait at least 14 days before attending another event



